Stirling's Formula and Laplace's Method
OR
How to Put Your Calculus to Good Use

The Undergraduate Colloquium
The University of Utah
December 4, 2001

Davar Khoshnevisan
Department of Mathematics
The University of Utah
davar@math.utah.edu
http://www.math.utah.edu/~davar

Jumbles (aka Permutations)

- ❄ There are 2 jumbles of "OF" ("OF" and "FO");

- ❄ There are 6 jumbles of "OFT" ("OFT", "FTO", "FOT", "OTF", "TOF", and "TFO");

- ❄ There are 24 jumbles of "SOFT" (!)

- ⋮

In general, there are $N! = N(N-1)\cdots 1$ jumbles of N distinct objects. This is read as "N factorial".

$N! = N \times (N-1) \times \cdots \times 1$ is a sequence that grows large rapidly as N grows, viz.,

$$
\begin{aligned}
1! &= 1 \\
2! &= 2 \\
3! &= 6 \\
4! &= 24 \\
5! &= 120 \\
&\vdots \\
20! &= 2.432902008 \times 10^{18} \\
40! &= 8.16 \times 10^{47} \\
100! &= 9.33 \times 10^{157} \\
400! &= \text{Error}
\end{aligned}
$$

$$\mathfrak{S}(N) = \sqrt{2\pi} e^{-N} N^{N+\frac{1}{2}}.$$

$\mathfrak{S}(20)$	$=$	$2.422786847 \times 10^{18}$
$\mathfrak{S}(40)$	$=$	$8.14217264483 \times 10^{47}$
$\mathfrak{S}(100)$	$=$	9.32×10^{157}
$\mathfrak{S}(400)$	$=$	Error

Stirling's Formula (De Moivre 1730)

$$\lim_{N\to\infty} \frac{N!}{\sqrt{2\pi}e^{-N}N^{N+\frac{1}{2}}} = 1.$$

In simple words, for large N,

$$N! \simeq \sqrt{2\pi}e^{-N}N^{N+\frac{1}{2}}.$$

A Little History

Stirling's formula was found by Abraham de Moivre and published in *"Miscellenea Analytica"* 1730. It was later refined, but published in the same year, by J. Stirling in *"Methodus Differentialis"* along with other little gems of thought. For instance, therein, Stirling computes the area under the *Bell Curve*:

$$\int_{-\infty}^{\infty} e^{-x^2/2}\, dx = \sqrt{2\pi},$$

$\cdots$ we will come back to this computation of James Stirling.

The Gamma Function Representation

$$N! = \int_0^\infty e^{-t} t^N \, dt.$$

To see this, let $u = t^N$ and $dv = e^{-t}\,dt$, and integrate by parts:

$$\begin{aligned}\int_0^\infty e^{-t} t^N \, dt &= -t^N e^{-t}\Big|_0^\infty + N \int_0^\infty e^{-t} t^{N-1} \, dt \\ &= N \int_0^\infty e^{-t} t^{N-1} \, dt.\end{aligned}$$

So, we can iterate this in N:

$$\begin{aligned}\int_0^\infty e^{-t} t^N \, dt &= N(N-1) \int_0^\infty e^{-t} t^{N-2} \, dt \\ &= N(N-1)(N-2) \int_0^\infty e^{-t} t^{N-3} \, dt \\ &\vdots \\ &= N! \int_0^\infty e^{-t} \, dt \\ &= N!.\end{aligned}$$

The Gamma Function

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1}\, dt, \qquad x > 1.$$

Check, by changing variables, that

$$\Gamma(\tfrac{1}{2}) = \int_0^\infty e^{-x^2/2}\, dx.$$

Stirling showed that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$. Thus, this is equivalent to the area under the bell curve $= \sqrt{2\pi}$ (check!)

Laplace's Method

Goal: estimate $\int_{-\infty}^{\infty} e^{Nf(x)}\,dx$ for large N, where f looks like



It can be shown that for such a function, all of the main contribution to the integral comes from the x values near x_0, say, $(1-\varepsilon)x_0 \leq x \leq (1+\varepsilon)x_0$. The contribution from the other x values is asymptotically (i.e., as $N \to \infty$) is negligible. Here, ε is an arbitrary positive constant.

So, for f of the form shown in the previous figure, and for large N,

$$\int_{-\infty}^{\infty} e^{Nf(x)}\,dx \simeq \int_{(1-\varepsilon)x_0}^{(1+\varepsilon)x_0} e^{Nf(x)}\,dx. \qquad (1)$$

Pick ε small and use Taylor's expansion near x_0:

$$f(x) \simeq f(x_0) + \frac{1}{2}f''(x_0)(x-x_0)^2,$$

and recall that $f''(x_0) < 0$ (Max. at x_0). Plug this in Eq. (1), to obtain

$$\int_{-\infty}^{\infty} e^{Nf(x)}\,dx \simeq e^{Nf(x_0)} \int_{(1-\varepsilon)x_0}^{(1+\varepsilon)x_0} e^{\frac{N}{2}f''(x_0)(x-x_0)^2}\,dx.$$

To re-iterate

$$\int_{-\infty}^{\infty} e^{Nf(x)}\,dx \simeq e^{Nf(x_0)} \int_{(1-\varepsilon)x_0}^{(1+\varepsilon)x_0} e^{\frac{N}{2}f''(x_0)(x-x_0)^2}\,dx.$$

Recall that $f''(x_0) < 0$, and change variables ($y = \sqrt{-Nf''(x_0)}(x - x_0)$):

$$\begin{aligned}\int_{-\infty}^{\infty} e^{Nf(x)}\,dx &\simeq e^{Nf(x_0)} \int_{-\varepsilon\sqrt{-Nf''(x_0)}}^{+\varepsilon\sqrt{-Nf''(x_0)}} e^{-\frac{1}{2}y^2}\,dy \times \\ &\quad \times \frac{1}{\sqrt{-Nf''(x_0)}} \\ &\simeq e^{Nf(x_0)} \int_{-\infty}^{\infty} e^{-\frac{1}{2}y^2}\,dy \times \\ &\quad \times \frac{1}{\sqrt{-Nf''(x_0)}} \\ &= \frac{\sqrt{2\pi}e^{Nf(x_0)}}{\sqrt{-Nf''(x_0)}}.\end{aligned}$$

Back to Stirling

Since $N! = \int_0^\infty e^{-t} t^N \, dt$ from the first slide,

$$N! = \int_0^\infty e^{-t + N \ln t} \, dt.$$

Change variables ($s = Nt$) to get

$$\begin{aligned} N! &= N^{N+1} \int_0^\infty e^{-N(\ln s - s)} \, ds \\ &= N^{N+1} \int_0^\infty e^{N f(s)} \, ds, \end{aligned}$$

where $f(s) = \ln(s) - s$. Check that f is of the desired form, $x_0 = 1$, $f(x_0) = -1$, $f''(x_0) = -1$, and now we derive the Striling's formula from Laplace's method easily:

$$N! \simeq N^{N+1} \frac{\sqrt{2\pi} e^{N f(x_0)}}{\sqrt{-N f''(x_0)}} = N^{N+1} \frac{\sqrt{2\pi} e^{-N}}{N^{\frac{1}{2}}},$$

which equals $\sqrt{2\pi} e^{-N} N^{N+\frac{1}{2}}$.

Easy Exercise

Check, by using Laplace's method, that

$$\int_0^\pi x^N \sin(x)\, dx \simeq \pi^{N+2} N^{-2}.$$

Harder Exercise

(Essentially the local de Moivre–Laplace central limit theorem)

Check, by using Laplace's method, that for even numbers N,

$$\frac{N!}{(N/2)!} \simeq 2^N \sqrt{\frac{2}{\pi N}},$$

where, in both exercises, $f(N) \simeq g(N)$ means that $f(N)/g(N) \to 1$, as $N \to \infty$.