


This article was downloaded by: [Universita Studi la Sapienza]

On: 29 April 2013, At: 02:03

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Molecular Physics: An International Journal at the Interface Between Chemistry and Physics

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tmph20>

Weighted density Lattice Boltzmann approach to fluids under confinement

Umberto Marini Bettolo Marconi ^a & Simone Melchionna ^b

^a Scuola di Scienze e Tecnologie, Università di Camerino, sez. di Perugia, Italy

^b CNR-IPCF, Consiglio Nazionale delle Ricerche, Dipartimento di Fisica, Università Sapienza, Roma, Italy

Accepted author version posted online: 14 Feb 2013. Version of record first published: 07 Mar 2013.

To cite this article: Umberto Marini Bettolo Marconi & Simone Melchionna (2013): Weighted density Lattice Boltzmann approach to fluids under confinement, *Molecular Physics: An International Journal at the Interface Between Chemistry and Physics*, DOI:10.1080/00268976.2013.771801

To link to this article: <http://dx.doi.org/10.1080/00268976.2013.771801>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

RESEARCH ARTICLE

Weighted density Lattice Boltzmann approach to fluids under confinement

Umberto Marini Bettolo Marconi^{a,*} and Simone Melchionna^b

^a*Scuola di Scienze e Tecnologie, Università di Camerino, Via Madonna delle Carceri, 62032 Camerino, INFN, sez. di Perugia, Italy;*

^b*CNR-IPCF, Consiglio Nazionale delle Ricerche, Dipartimento di Fisica, Università Sapienza, Piazzale A. Moro 2, 00185 Roma, Italy*

(Received 2 December 2012; final version received 25 January 2013)

The Enskog-like kinetic approach, recently introduced by us to study strongly inhomogeneous fluids, is reconsidered in order to improve the description of the transport coefficients. The approach is based on a separation of the interaction between hydrodynamic and non-hydrodynamic parts. The latter is treated within a simple relaxation approximation. We show that, by considering the non-hydrodynamic part via a weighted density approximation, we obtain a better prediction of the transport coefficients. By virtue of the simplicity of the kinetic equation we are able to solve numerically the phase space distribution in the presence of inhomogeneities, such as confining surfaces, via a Lattice Boltzmann method. Analytical estimates of the importance of these corrections to the transport coefficients in bulk conditions is provided. Poiseuille flow of the hard-sphere fluid confined between two parallel smooth walls is studied and their pore-averaged properties are determined.

Keywords: nanofluidics; Lattice Boltzmann method; density functional theory; weighted density approximation; viscosity

1. Introduction

The technological and industrial relevance of processes where molecular fluids flow through channels of nanometric diameter require to develop methods to control and predict such phenomena [1–6]. Among the various techniques available today, one can mention continuum-based computational fluid dynamics methods, finite-element methods for Navier–Stokes equations [7], molecular dynamics [8,9], kinetic theory, [10] dynamic Boltzmann free-energy functional theory [11] and dynamical density functional theory for dense atomic liquids [12,13]. In these methods, one studies either the evolution of the local particle density $n(\mathbf{r}, t)$ or the evolution of the one-particle phase-space density distribution function.

The strategy based on the solution of the phase-space distribution function is known as the Lattice Boltzmann method [14,15] and is based on a discretised version of the Boltzmann transport equation, originally conceived to solve at a reduced computational cost the fluid dynamics equations. It can deal with complicated geometries and is numerically robust but, conversely, is typically based on top-down strategy: the phenomenological parameters, such as the viscosity and the thermal conductivity, are utilised as lumped parameter whose values are provided by some experiment or macroscopic model. Since the underlying microscopic level is unexplored, in principle one cannot infer the nature of the atomic constituents. In addition, since the phenomenological transport coefficients are fixed from bulk

measurements, it is awkward to control how they change under inhomogeneous conditions, such as those due to confinement in narrow spaces.

In a series of papers, we have considered an alternative approach [16–18], that does not require the *a priori* knowledge of the transport coefficients, nor the equation of state, but directly tackles the Boltzmann equation corresponding to a well defined microscopic model. A convenient route is to consider a system of hard-spheres and possibly adding attractive potential tails. The Hamiltonian determines the cross-section entering the Boltzmann–Enskog (BE) equation, thus establishing a link between microscopic and macroscopic properties. Using such bottom-up approach, it is possible to determine all transport coefficients and the equation of state from the BE equation, without introducing empirical ingredients. Such a strategy was pioneered by Luo, who carried out a systematic derivation of the Lattice Boltzmann equation starting from the Enskog equation. He obtained the equation of state for non-ideal gases together with thermodynamic consistency [19].

The present method not only takes into account the non-local nature of the collision operator and leads to a non-ideal gas equation of state, but also reproduces the dependence of the transport coefficients on the density. The approximation scheme utilises concepts inspired by the density functional theory in the weighted density approximation (WDA) version [20–21], as regards to the use of a smoothed density field and the equilibrium pair distribution function as

*Corresponding author. Email: umberto.marinibettolo@unicam.it

key quantities. At equilibrium our approach gives the same equations of the WDA and for such a reason we name it weighted density Lattice Boltzmann method (WDLB). Finally, the method is employed to derive a practical numerical scheme to study liquids flowing in confined geometries.

The purpose of this article is also to improve the WDLB by considering corrections to the viscosity and to the thermal conductivity. Such corrections are already present in the Enskog theory and, since they arise from the coupling between kinetic and collisional modes, they are named cross-over contributions to the transport coefficients and depend linearly with the fluid density. To this aim, we follow the clever approximation introduced by Dufty, Santos and Brey (DSB) [22] to simplify the revised Enskog theory (RET) equation [23], without spoiling its good features. In particular, this approach preserves the non-locality of interactions, it accounts for the static pair correlations in a realistic way and provides a reasonable equation of state. In essence, Brey and co-workers proposed to replace the complex collision kernel featuring in the original RET (see Equation (24) below) by a more tractable object.

The idea behind this proposal is to separate the relevant slow hydrodynamic modes from the fast evolving and less relevant kinetic modes. The first ones are then carefully handled, whereas the latter are treated by resorting to a simple relaxation time approximation, thus avoiding the effort of computing difficult integrals involving high velocity momenta of the collision operator. The resulting approximated collision operator consists of two pieces, one acting on the hydrodynamic modes and the second determining the relaxation towards equilibrium of the kinetic modes through the simplest approximation of the Boltzmann equation, the Bhatnagar-Gross-Krook (BGK) kernel [24].

Through this simplification, one obtains a representation where a set of effective fields describes the effect of intermolecular interactions in a self-consistent fashion, and the resulting transport equation is amenable to numerical solution. In a series of papers, [25–27] we have numerically implemented this approach [22] by developing a numerical algorithm in the framework of the Lattice Boltzmann method. Here, we consider the effect of a more refined treatment of the relaxation term, as proposed by Santos, Montanero, Dufty and Brey (SMDB) [28] in order to account for the cross-over contributions to the transport coefficients.

This paper is organised as follows. In Section 2, we briefly review the employed kinetic method in order to present the extension of the WDLB approach, and motivate the introduction of the new terms into the equation of evolution used in numerical applications. In Section 3, we give the explicit representation of the collision operator in the case of hard spheres and compute the effective fields. In Section 4, we present results for the transport coefficients. Finally, in Section 5, we present the numerical test of the

new approximation. In Section 6, we make some concluding remarks.

2. Model

The evolution of the phase-space one-particle distribution $f(\mathbf{r}, \mathbf{v}, t)$ is represented by the following transport equation [29–31]:

$$\partial_t f(\mathbf{r}, \mathbf{v}, t) + v_i \partial_i f(\mathbf{r}, \mathbf{v}, t) + \frac{F_i(\mathbf{r})}{m} \frac{\partial}{\partial v_i} f(\mathbf{r}, \mathbf{v}, t) = \Omega[f](\mathbf{r}, \mathbf{v}, t), \quad (1)$$

where \mathbf{F} is an external force and $\Omega[f]$ represents the effect of the interactions among the fluid particles which we shall specify later. We adopt the Einstein summation convention that repeated indices are implicitly summed over and the notation ∂_i to indicate the partial derivative with respect to the i th component of the vector \mathbf{r} and ∂_t to indicate the partial derivative with respect to time.

In principle, after giving an explicit representation of Ω according to the underlying microscopic model, Equation (1) can be solved numerically. The case of non-local functionals Ω , such as the RET, is computationally demanding. Alternatively, one can consider very simple forms of Ω , the BGK being the simplest *ad hoc* choice. Our strategy considers a non-local functional Ω , but factorises the velocity dependence in terms of a local Maxwellian distribution times Hermite tensor polynomials [25]. In addition, since one is chiefly interested in the evolution of the hydrodynamic variables it is possible to further simplify the form of Ω without altering the local transfer of momentum and energy.

In this framework, we implicitly assume that after few molecular collisions the system reaches a state of local thermodynamic equilibrium characterised by the hydrodynamic variables $n(\mathbf{r}, t)$, $\mathbf{u}(\mathbf{r}, t)$, $T(\mathbf{r}, t)$, being the number density, local fluid velocity and local temperature, respectively. We further assume that the phase space distribution depends on space and time only through these fields and their gradients.

The five hydrodynamic fields can be obtained from $f(\mathbf{r}, \mathbf{v}, t)$ through the relations

$$\begin{pmatrix} n(\mathbf{r}, t) \\ n(\mathbf{r}, t)\mathbf{u}(\mathbf{r}, t) \\ \frac{3}{2}k_B n(\mathbf{r}, t)T(\mathbf{r}, t) \end{pmatrix} = \int d\mathbf{v} \begin{pmatrix} 1 \\ \mathbf{v} \\ \frac{m(\mathbf{v}-\mathbf{u}(\mathbf{r}, t))^2}{2} \end{pmatrix} f(\mathbf{r}, \mathbf{v}, t), \quad (2)$$

where k_B is the Boltzmann constant.

We also introduce the corresponding moments of the collision operator

$$\begin{pmatrix} 0 \\ \mathbf{C}(\mathbf{r}, t) \\ Q(\mathbf{r}, t) \end{pmatrix} = \int d\mathbf{v} \begin{pmatrix} 1 \\ m\mathbf{v} \\ \frac{m(\mathbf{v}-\mathbf{u}(\mathbf{r}, t))^2}{2} \end{pmatrix} \Omega(\mathbf{r}, \mathbf{v}, t). \quad (3)$$

By multiplying both sides of Equation (1) by the lowest Hermite tensorial polynomials (that is, 1 , \mathbf{v} , and $m(\mathbf{v} - \mathbf{u})^2/2$) and integrating with respect to \mathbf{v} , we obtain the following set of balance equations for the five hydrodynamic variables:

$$\partial_t n(\mathbf{r}, t) + \partial_i (n(\mathbf{r}, t) u_i(\mathbf{r}, t)) = 0, \quad (4)$$

$$mn(\mathbf{r}, t)[\partial_t u_j(\mathbf{r}, t) + u_i(\mathbf{r}, t)\partial_i u_j(\mathbf{r}, t)] + \partial_i \tilde{P}_{ij}(\mathbf{r}, t) - F_j(\mathbf{r})n(\mathbf{r}, t) - C_j(\mathbf{r}, t) = 0 \quad (5)$$

and

$$\frac{3}{2}n(\mathbf{r}, t)k_B[\partial_t T(\mathbf{r}, t) + u_i(\mathbf{r}, t)\partial_i T(\mathbf{r}, t)] + \tilde{P}_{ij}(\mathbf{r}, t)\partial_i u_j(\mathbf{r}, t) + \partial_i \tilde{q}_i(\mathbf{r}, t) - Q(\mathbf{r}, t) = 0. \quad (6)$$

The momentum and temperature Equations (5) and (6) contain two additional quantities, a pressure term

$$\tilde{P}_{ij}(\mathbf{r}, t) = m \int d\mathbf{v} f(\mathbf{r}, \mathbf{v}, t)(\mathbf{v} - \mathbf{u})_i(\mathbf{v} - \mathbf{u})_j \quad (7)$$

and a heat flux term:

$$\tilde{q}_i(\mathbf{r}, t) = \frac{m}{2} \int d\mathbf{v} f(\mathbf{r}, \mathbf{v}, t)(\mathbf{v} - \mathbf{u})^2(\mathbf{v} - \mathbf{u})_i, \quad (8)$$

which in the $n \rightarrow 0$ limit can be identified with the kinetic components of the pressure tensor and heat flux, respectively. As we shall see in Section 3, these two quantities also contain contributions which are neither purely kinetic nor purely collisional.

One can also identify the last term in Equation (5) with the gradient of the collisional contribution to the pressure tensor, P_{ji}^c ,

$$C_i(\mathbf{r}, t) = -\frac{1}{m} \partial_j P_{ji}^c(\mathbf{r}, t) \quad (9)$$

and Q with a combination of the gradient of the collisional contribution to the heat flux vector q_i^c and the pressure tensor times the strain rate:

$$Q(\mathbf{r}, t) = -\frac{1}{m} [\partial_j q_j^c(\mathbf{r}, t) + P_{ji}^c(\mathbf{r}, t)\partial_j u_i(\mathbf{r}, t)]. \quad (10)$$

The second equality in Equations (9) and (10) identifies the effective fields C_i and B with the thermodynamic fields.

Formally, we have derived a set of equations for the five hydrodynamic variables, but these do not form a closed system, because we still need either to specify the non-equilibrium distribution function f or to assume some phenomenological constitutive relations between the gradients of P_{ij} and q (the sums of the kinetic and collisional parts)

and the fluxes. Using the latter strategy one obtains the Navier-Stokes equations [32].

The alternative route is to focus on the evolution of f , with the collision operator Ω being in general a nonlinear functional of f , so that any kind of analytic work results hard. The simplest and crudest way to proceed is to approximate Ω by the following relaxation ansatz as done by BGK [24]

$$\Omega^{BGK}(\mathbf{r}, \mathbf{v}, t) = -\omega_0 [f(\mathbf{r}, \mathbf{v}, t) - n(\mathbf{r}, t)\phi_M(\mathbf{r}, \mathbf{v}, t)], \quad (11)$$

where

$$\phi_M(\mathbf{r}, \mathbf{v}, t) = \left[\frac{m}{2\pi k_B T(\mathbf{r}, t)} \right]^{3/2} \exp\left(-\frac{m(\mathbf{v} - \mathbf{u})^2}{2k_B T(\mathbf{r}, t)}\right)$$

is the local Maxwellian. The BGK recipe for Ω preserves the number of particles, the momentum and the kinetic energy, in other words it fulfills the physical symmetries and conservation laws of the fluid. Finally, it gives a quite uninteresting thermodynamic behaviour corresponding to an ideal gas. Instead, it gives non-vanishing values of the transport coefficients, which turn out to be functions of phenomenological parameter ω_0 [24].

In view of the interest for the non-ideal gas behaviour, one seek for a better approximation capable of predicting a non-trivial equation of state and transport coefficients. The Boltzmann equation for hard spheres has non-trivial transport coefficients but lacks a satisfactory equation of state. The RET form of Ω has the sought features, but is hardly tractable in numerical work [23].

An interesting and practical reduction of the RET collision operator has been devised by Dufty *et al.* [22]. Their idea is to separate the contributions of Ω to the hydrodynamic equations from those affecting the evolution of the non-hydrodynamic modes, by projecting the collision term onto the hydrodynamic subspace spanned by the functions $\{1, \mathbf{v}, v^2\}$ and onto the complementary kinetic subspace:

$$\Omega = \mathcal{P}_{\text{hydro}}\Omega + (I - \mathcal{P}_{\text{hydro}})\Omega \quad (12)$$

with

$$\mathcal{P}_{\text{hydro}}\Omega = \frac{1}{k_B T(\mathbf{r}, t)} \phi_M(\mathbf{r}, t) \left[(\mathbf{v} - \mathbf{u}) \cdot \mathbf{C}(\mathbf{r}, t) + \left[\frac{m(\mathbf{v} - \mathbf{u}(\mathbf{r}, t))^2}{3k_B T} - 1 \right] Q(\mathbf{r}, t) \right]. \quad (13)$$

The second term, $\Omega_{\perp} \equiv (I - \mathcal{P}_{\text{hydro}})\Omega$ is further approximated as:

$$\Omega_{\perp} \approx -\omega_0 [f(\mathbf{r}, \mathbf{v}, t) - n(\mathbf{r}, t)\phi_M(\mathbf{r}, \mathbf{v}, t)] \quad (14)$$

and Equation (1) by a simpler equation, where the complicated interaction between non-hydrodynamic modes is approximated via a BGK-like relaxation term with ω_0 being a phenomenological collision frequency chosen as to reproduce the kinetic contribution to the viscosity.

It was pointed out by Santos et al. [28] that better results for the transport properties could be obtained upon modifying the BGK relaxation term in Equation (14) as to allow the relaxation time to depend on the spatial inhomogeneities of the local equilibrium state. This effect depends on the collisional transfer mechanism associated with the difference in position of the colliding particles. To achieve this goal, following Santos, Montanero, Dufty and Brey [28], we complement the relaxation term in Equation (14) in the following way:

$$\begin{aligned} \Omega_{\perp}^{SMDB}(\mathbf{r}, t) = & -\omega_0[f(\mathbf{r}, \mathbf{v}, t) - n(\mathbf{r}, t)\phi_M(\mathbf{r}, \mathbf{v}, t)] + (m\beta)^2\phi_M(\mathbf{r}, \mathbf{v}, t) \\ & \times \left\{ \left[(\mathbf{v} - \mathbf{u})_i(\mathbf{v} - \mathbf{u})_j - \frac{1}{3}(\mathbf{v} - \mathbf{u})^2\delta_{ij} \right] \frac{1}{2}A_{ij}(\mathbf{r}, t) \right. \\ & \left. + \left[\frac{m\beta(\mathbf{v} - \mathbf{u})^2}{2} - \frac{5}{2} \right] (\mathbf{v} - \mathbf{u})_i \frac{1}{5}B_i(\mathbf{r}, t) \right\}. \end{aligned} \quad (15)$$

The extra terms in Equation (15) give a better representation of the decays towards equilibrium as discussed in Ref. [28]. The first term is proportional to ω_0 , which assumes a constant value throughout the system, whereas the second term may depend on the spatial inhomogeneities of the velocity and of the temperature fields. Such a dependence occurs through the functions $A_{ij}(\mathbf{r}, t)$ and $B_i(\mathbf{r}, t)$ and thus is modulated by the spatial structure of the fluid. In the case of a uniform system, the relaxation process may occur only via the term proportional to ω_0 , when the distribution function f is different from the local equilibrium distribution. The inclusion of these terms also yields a quantitative improvement of the transport coefficients with respect the simpler theory discussed previously.

The new terms can be obtained by computing the following projections of the collision operator:

$$\begin{aligned} A_{ij}(\mathbf{r}, t) = \int d\mathbf{v} \left[(\mathbf{v} - \mathbf{u})_i(\mathbf{v} - \mathbf{u})_j \right. \\ \left. - \frac{1}{3}(\mathbf{v} - \mathbf{u})^2\delta_{ij} \right] \Omega(\mathbf{v}, \mathbf{r}, t) \end{aligned} \quad (16)$$

and

$$B_i(\mathbf{r}, t) = \int d\mathbf{v} \left[(\mathbf{v} - \mathbf{u})^2 - 5\frac{k_B T}{m} \right] (\mathbf{v} - \mathbf{u})_i \Omega(\mathbf{v}, \mathbf{r}, t) \quad (17)$$

By construction, the new terms do not affect the structure of the continuity Equation (4), momentum balance Equation (5) and energy balance Equation (6). The SMDB

equation can be cast in the form:

$$\begin{aligned} \partial_t f(\mathbf{r}, \mathbf{v}, t) + v_i \partial_i f(\mathbf{r}, \mathbf{v}, t) \\ + \frac{F_i(\mathbf{r})}{m} \frac{\partial}{\partial v_i} f(\mathbf{r}, \mathbf{v}, t) = \Omega_{\perp}^{SMDB}(\mathbf{r}, t) \\ + m\beta\phi_M(\mathbf{r}, \mathbf{v}, t) \left\{ (\mathbf{v} - \mathbf{u}) \cdot \mathbf{C}(\mathbf{r}, t) \right. \\ \left. + \left(\frac{m\beta(\mathbf{v} - \mathbf{u})^2}{3} - 1 \right) Q(\mathbf{r}, t) \right\}, \end{aligned} \quad (18)$$

where $\beta = 1/k_B T(\mathbf{r}, t)$. A simple calculation allows to compute the deviation δf of the distribution function $f(\mathbf{r}, \mathbf{v}, t)$ from its local equilibrium value $f_{loc} = n(\mathbf{r}, t)\phi_M(\mathbf{r}, t)$. We replace $f \rightarrow f_{loc}$ in the left-hand side of Equation (18) and obtain:

$$\begin{aligned} \left\{ \left[\partial_t n + \partial_i(nu_i) \right] + \left[n\partial_t u_k + nu_i \partial_i(nu_k) + \partial_k(nT) \right. \right. \\ \left. - \frac{F_k}{m} - C_k \right] \frac{m(v_k - u_k)}{k_B T} + \frac{n}{2T} \left[\partial_i T + u_i \partial_i T \right. \\ \left. + \frac{2}{3} T \partial_i u_i - \frac{2}{3n} Q \right] \left(\frac{m(\mathbf{v} - \mathbf{u})^2}{k_B T} - 3 \right) \\ \left. + \left[\frac{n}{T} \partial_i T - \frac{m\beta}{5} B_i \right] \left(m \frac{(\mathbf{v} - \mathbf{u})^2}{2k_B T} - \frac{5}{2} \right) (v_i - u_i) \right. \\ \left. + m\beta \left[n \partial_i u_k - \frac{m\beta}{2} A_{ik} \right] \left((v_i - u_i)(v_k - u_k) \right. \right. \\ \left. \left. - \frac{(\mathbf{v} - \mathbf{u})^2}{3} \delta_{ij} \right) \right\} \phi_M(\mathbf{r}, \mathbf{v}, t) = -\omega_0 \delta f(\mathbf{r}, \mathbf{v}, t). \end{aligned} \quad (19)$$

Since the projections of δf over $(1, (\mathbf{v}_i - u_i), m(\mathbf{v} - \mathbf{u})^2/k_B T - 3)$ vanish, we impose that also the first three terms in the left-hand side vanish. These are the so-called solvability conditions and are precisely the balance Equations (4)–(6). We thus obtain the following explicit representation of δf

$$\begin{aligned} \delta f = -\frac{1}{\omega_0} \phi_M(\mathbf{r}, \mathbf{v}, t) \left\{ \left[\frac{n(\mathbf{r}, t)}{T} \partial_i T(\mathbf{r}, t) \right. \right. \\ \left. - \frac{(m\beta)^2}{5} B_i \right] \left(\frac{m(\mathbf{v} - \mathbf{u})^2}{2k_B T(\mathbf{r}, t)} - \frac{5}{2} \right) (v_i - u_i) \\ \left. + m\beta \left[n(\mathbf{r}, t) \partial_i u_k(\mathbf{r}, t) - \frac{m\beta}{2} A_{ik} \right] \right. \\ \left. \times \left((v_i - u_i)(v_k - u_k) - \frac{(\mathbf{v} - \mathbf{u})^2}{3} \delta_{ik} \right) \right\}. \end{aligned} \quad (20)$$

The fields A_{ij} and B_i represent a measure of the distortion of $f(\mathbf{r}, \mathbf{v}, t)$ with respect to the local equilibrium $f_{loc}(\mathbf{r}, \mathbf{v}, t)$. Using such an approximation for δf , we can now estimate

how the distortion affects the components of the pressure tensor (7):

$$\begin{aligned} \tilde{P}_{ij}(\mathbf{r}, t) = k_B T(\mathbf{r}, t) n(\mathbf{r}, t) & \left\{ \delta_{ij} - \frac{1}{\omega_0} \left((\partial_i u_j(\mathbf{r}, t) \right. \right. \\ & \left. \left. + \partial_j u_i(\mathbf{r}, t)) - \frac{2}{3} \delta_{ij} \sum_k \partial_k u_k(\mathbf{r}, t) \right) \right\} + \delta P_{ij}(\mathbf{r}, t), \end{aligned} \quad (21)$$

where the first term is the BGK result and

$$\delta P_{ij}(\mathbf{r}, t) = \frac{m}{\omega_0} \left[A_{ij}(\mathbf{r}, t) + A_{ji}(\mathbf{r}, t) - \frac{2}{3} \delta_{ij} \sum_k A_{kk}(\mathbf{r}, t) \right] \quad (22)$$

stems from the A_{ij} contribution. The heat flux is obtained by substituting δf in Equation (8)

$$\tilde{q}_i(\mathbf{r}, t) = -\frac{5}{2} \frac{1}{m\omega_0} n(\mathbf{r}, t) k_B^2 T(\mathbf{r}, t) \partial_i T(\mathbf{r}, t) + \delta q_i(\mathbf{r}, t) \quad (23)$$

with $\delta q_i(\mathbf{r}, t) = \frac{1}{\omega_0} \frac{m}{2} B_i$.

It is worth noting that Luo [33] compared the approaches derived from the Enskog equation with those based on the simplifying assumption that the intermolecular forces are assimilated to a forcing self-consistent term. He concluded that the latter are plagued by an inconsistency. In fact, the pressure featuring in the momentum equation must also feature in the energy equation, a condition which is not satisfied if one treats the interactions in a mean-field fashion [34,35]. In this respect, the present treatment has the correct form, since Equation (6) contains the viscous dissipation term representing the irreversible transformation of kinetic energy into heat.

3. Hard sphere model

At this stage, we are interested in obtaining an explicit representation of the fields (C_i, Q, A_{ij}, B_i) in order to determine the viscosity and thermal conductivity, and to implement the numerical solution of the evolution Equation (18) by lattice methods. To this purpose, we use as a benchmark the prototypical model of dense fluids, namely the hard-sphere system at moderate packing fractions. The RET collision operator involves only the single-particle phase space distribution and accounts for static pair correlations, but not velocity correlations. At high density, it becomes more inaccurate because it disregards the occurrence of correlated collisions. Nevertheless, the RET approach is very accurate for our purposes and has the interesting feature of taking into account the instantaneous transfer of energy and momentum between particles whose distance equals the hard

sphere diameter. For this reason, even to a spatially uniform state corresponds a non-ideal gas pressure.

The RET collision operator reads [23]:

$$\begin{aligned} \Omega^{RET}(\mathbf{r}, \mathbf{v}_1, t) = \sigma^{d-1} \int d\mathbf{v}_2 \int d\hat{\mathbf{k}} \Theta(\hat{\mathbf{k}} \cdot \mathbf{v}_{12}) (\hat{\mathbf{k}} \cdot \mathbf{v}_{12}) \\ \times \{ g_2(\mathbf{r}, \mathbf{r} - \hat{\mathbf{k}}\sigma, t) f(\mathbf{r}, \mathbf{v}'_1, t) f(\mathbf{r} - \hat{\mathbf{k}}\sigma, \mathbf{v}'_2, t) \\ - g_2(\mathbf{r}, \mathbf{r} + \hat{\mathbf{k}}\sigma, t) f(\mathbf{r}, \mathbf{v}_1, t) f(\mathbf{r} + \hat{\mathbf{k}}\sigma, \mathbf{v}_2, t) \}, \end{aligned} \quad (24)$$

where \mathbf{v}'_1 and \mathbf{v}'_2 are scattered velocities determined from $\mathbf{v}'_1 = \mathbf{v}_1 - (\hat{\mathbf{k}} \cdot \mathbf{v}_{12}) \hat{\mathbf{k}}$ and $\mathbf{v}'_2 = \mathbf{v}_2 + (\hat{\mathbf{k}} \cdot \mathbf{v}_{12}) \hat{\mathbf{k}}$, σ is the hard-sphere diameter, $\hat{\mathbf{k}}$ is the unit vector directed from one particle to another, Θ is the Heaviside function and $g_2(\mathbf{r}_1, \mathbf{r}_2, t)$ is the pair correlation function evaluated at contact distance $|\mathbf{r} - \mathbf{r}'| = \sigma$. It describes the positional pair correlations corresponding to the inhomogeneous single particle density profile $n(\mathbf{r}, t)$. There is no velocity dependence in the two particle correlation.

Following Ref. [36], we assume that g_2 at time t is the same as the one corresponding to the same system having a density profile equal to $n(\mathbf{r}, t)$ at thermodynamic equilibrium, an assumption often referred to as the adiabatic approximation. Moreover, since the exact inhomogeneous form of g_2 is unknown, we resort to the Fischer–Methfessel prescription [37], stating that $g_2(\mathbf{r}_1, \mathbf{r}_2, t)$ is given by the pair correlation function at contact of an homogeneous system evaluated replacing n by the following smeared density:

$$\bar{n}(\mathbf{r}, t) \equiv \frac{6}{\pi \sigma^3} \int_{|\mathbf{r}'| < \sigma/2} d\mathbf{r}' n(\mathbf{r} + \mathbf{r}'). \quad (25)$$

Because we have access only to the first lower velocity moments of f , we cannot solve the full problem (1) using such a collision operator. The DSB and SMDB recipes represent viable options. Recently, we have used the SDB equation in conjunction with the Lattice Boltzmann method to study one component, two component and ternary mixtures with interesting applications also to electrolyte solutions [38,39]. Hereafter, we want to test the SMDB equation implemented by an appropriate LB method and compare the results.

The expressions of \mathbf{C}, B have been obtained before [17] and read

$$\begin{aligned} C_i(\mathbf{r}, t) = -k_B T(\mathbf{r}, t) n(\mathbf{r}, t) \sigma^2 \int d\hat{\mathbf{k}} k_i g_2 \\ \times (\mathbf{r}, \mathbf{r} + \sigma \hat{\mathbf{k}}, t | n(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) \left[1 - \frac{2}{\sqrt{\pi} k_B T(\mathbf{r}, t) / m} \hat{\mathbf{k}} \right. \\ \left. \cdot [\mathbf{u}(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - \mathbf{u}(\mathbf{r}, t)] + \frac{T(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - T(\mathbf{r}, t)}{2T(\mathbf{r}, t)} \right] \end{aligned} \quad (26)$$

and

$$\begin{aligned}
Q(\mathbf{r}, t) &= k_B T(\mathbf{r}, t) n(\mathbf{r}, t) \sigma^2 \int d\hat{\mathbf{k}} g_2(\mathbf{r}, \mathbf{r}) \\
&+ \sigma \hat{\mathbf{k}} \cdot t |n(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) \left[-\frac{\hat{\mathbf{k}} \cdot [\mathbf{u}(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - \mathbf{u}(\mathbf{r}, t)]}{2} \right. \\
&+ \left. \frac{1}{\sqrt{\pi}} \sqrt{\frac{k_B T(\mathbf{r}, t)}{m}} \frac{T(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - T(\mathbf{r}, t)}{T(\mathbf{r}, t)} \right]. \quad (27)
\end{aligned}$$

A similar calculation was performed to evaluate A_{ij} and B_i using Equations (16) and (17), respectively, under the approximation that $\Omega^{RET}[f, f]$ featuring in these two equations was replaced by $\Omega^{RET}[f_{loc}, f_{loc}]$ [13–40]. We find:

$$\begin{aligned}
A_{ij}(\mathbf{r}, t) &= -\frac{k_B T(\mathbf{r}, t)}{5m} n(\mathbf{r}, t) \sigma^2 \int d\hat{\mathbf{k}} g_2(\mathbf{r}, \mathbf{r}) \\
&+ \sigma \hat{\mathbf{k}} \cdot t |n(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) \left[\hat{k}_j [u_i(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - u_i(\mathbf{r}, t)] \right. \\
&+ \left. \hat{k}_i [u_j(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - u_j(\mathbf{r}, t)] \right]. \quad (28)
\end{aligned}$$

Similarly, we derive the correction to the heat flux vector

$$\begin{aligned}
B_i(\mathbf{r}, t) &= -2 \frac{k_B^2}{m^2} T(\mathbf{r}, t) n(\mathbf{r}, t) \sigma^2 \int d\hat{\mathbf{k}} \hat{k}_i g_2 \\
&\times (\mathbf{r}, \mathbf{r} + \sigma \hat{\mathbf{k}} \cdot t |n(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) \left[T(\mathbf{r} + \sigma \hat{\mathbf{k}}, t) - T(\mathbf{r}, t) \right]. \quad (29)
\end{aligned}$$

As shown in Ref. [26], a useful representation of the term (26) is obtained by decomposing it in the following way: $\mathbf{C}(\mathbf{r}, t) = n(\mathbf{r}, t) (\mathbf{F}^{mf}(\mathbf{r}, t) + \mathbf{F}^{viscous}(\mathbf{r}, t))$, where we can identify a force acting on the particle at position \mathbf{r} as due to the influence of all remaining particles in the system, the so-called potential of mean force [41]:

$$F_i^{mf}(\mathbf{r}, t) = -k_B T \sigma^2 \int d\hat{\mathbf{k}} \hat{k}_i g_2(\mathbf{r}, \mathbf{r} + \sigma \mathbf{k}, t) n(\mathbf{r} + \sigma \mathbf{k}, t) \quad (30)$$

and a viscous term

$$\begin{aligned}
F_i^{viscous}(\mathbf{r}, t) &= 2\sigma^2 \sqrt{\frac{mk_B T}{\pi}} \int d\hat{\mathbf{k}} \hat{k}_i k_j g_2 \\
&\times (\mathbf{r}, \mathbf{r} + \sigma \mathbf{k}, t) n(\mathbf{r} + \sigma \mathbf{k}, t) (u_j(\mathbf{r} + \sigma \mathbf{k}) - u_j(\mathbf{r})) \quad (31)
\end{aligned}$$

and a force due to the presence of thermal gradients

$$\begin{aligned}
F_i^T(\mathbf{r}, t) &= -\frac{\sigma^2}{2} \int d\hat{\mathbf{k}} \hat{k}_i g_2(\mathbf{r}, \mathbf{r} + \sigma \mathbf{k}, t) n \\
&\times (\mathbf{r} + \sigma \mathbf{k}, t) k_B [T(\mathbf{r} + \sigma \mathbf{k}, t) - T(\mathbf{r}, t)]. \quad (32)
\end{aligned}$$

One can see that the form of the fields C_i , Q , A_{ij} , B_i depends on density, velocity profile and temperature. Only one contribution to C_i inherits the property of the density functional theory of being fully determined by the density only. The other terms require the knowledge of the hydrodynamic fields even in the present approximation, since they are not slaved to the density field as it occurs in dynamical density functional theory.

4. Analytic estimate of the bulk transport coefficients from WDLB

Let us consider the limit of small velocity gradients and constant density and temperature, so that:

$$A_{ij}(\mathbf{r}, t) \simeq -\frac{k_B T}{m} n^2 g_2(\sigma^+) \frac{4\pi}{15} \sigma^3 [\partial_j u_i + \partial_i u_j] \quad (33)$$

and, using Equation (22):

$$\begin{aligned}
\delta P_{ij}(\mathbf{r}, t) &= -\frac{1}{\omega_0} k_B T n^2 g_2(\sigma^+) \frac{8\pi}{15} \sigma^3 \\
&\times \left[\partial_j u_i + \partial_i u_j - \frac{2}{3} \delta_{ij} \sum_k \partial_k u_k \right]. \quad (34)
\end{aligned}$$

Let us consider a system of uniform density and temperature, and a small sinusoidal velocity perturbation varying in a direction normal to the streamlines:

$$\mathbf{u}(x, t) = (0, u_y(0, t) e^{iq_x x}, 0). \quad (35)$$

By neglecting the nonlinear term, we obtain from Equation (5)

$$\frac{\partial}{\partial t} u_y(x, t) = -\frac{\partial}{\partial x} \tilde{P}_{xy}(x, t) + C_y(x, t) \quad (36)$$

and, substituting the expressions for C_y and \tilde{P}_{xy} ,

$$\begin{aligned}
n \frac{\partial}{\partial t} u_y(x, t) &= \left[\frac{1}{\omega_0} k_B T n \left(1 + \frac{8\pi}{15} n \sigma^3 g_2(\sigma^+) \right) \right. \\
&+ \left. \frac{4}{15} \sqrt{m\pi k_B T} n \sigma^4 g_2(\sigma^+) \right] \frac{\partial^2 u_y(x, t)}{\partial x^2}. \quad (37)
\end{aligned}$$

The solutions have exponential form, $u_y(x, t) = u_y(0, t) e^{iq_x x} e^{-(\eta/n)q_x^2 t}$, where the shear viscosity η is given by the expression inside square brackets in the right-hand side of Equation (37).

The first term arises from the reduction of the velocity gradients due to the displacement of particles from a region characterised by a higher fluid velocity \mathbf{u} to a region of lower velocity and vice versa. The third term takes into

account the fact that neighbouring volume elements moving at different velocities can exchange momentum due to the interaction of their surface molecules without involving particle displacement. The second term $\eta^{(1)}$, called cross-over term, has a mixed nature [42]. Finally, we set the collision frequency to the value $\omega_0 = \frac{16}{5}n\sigma^2 g_2(\sigma)\sqrt{\pi k_B T/m}$ [28] giving the correct low density shear viscosity of hard spheres of diameter σ

$$\eta = \frac{nk_B T}{\omega_0} \left[1 + \frac{8\pi}{15}n\sigma^3 g_2(\sigma^+) \right] + \frac{4}{15}\sqrt{m\pi k_B T}n\sigma^4 g_2(\sigma^+). \quad (38)$$

4.1. Bulk viscosity

In order to derive the expression for the bulk viscosity, η_b , we consider a velocity field of the form $\mathbf{u} = (u_x(x, t), 0, 0)$, use the macroscopic relation

$$\frac{\partial P_{xx}(x, t)}{\partial x} = -\left(\eta_b + \frac{4}{3}\eta\right) \frac{\partial^2 u_x(x, t)}{\partial x^2} \quad (39)$$

to define η_b and obtain after a simple calculation

$$\frac{\partial P_{xx}(x, t)}{\partial x} = -\left[\frac{4}{3} \frac{1}{\omega_0} k_B T n \left(1 + \frac{8\pi}{15} n \sigma^3 g_2(\sigma^+) \right) + \frac{4}{5} \sqrt{\pi} \sqrt{m k_B T} n \sigma^4 g_2(\sigma^+) \right] \frac{\partial^2 u_x(x, t)}{\partial x^2}. \quad (40)$$

By comparing, the two expressions for the derivative of the pressure tensor, we find $\eta_b = \frac{5}{3}\eta$ and conclude that η_b is purely collisional with no kinetic and cross contributions.

4.2. Thermal conductivity

The heat flux correction B_i for small gradients can be approximated as

$$\delta \tilde{q}_x(\mathbf{r}, t) \simeq -\frac{1}{\omega_0} \frac{k_B^2}{m} T(\mathbf{r}, t) n^2 g_2(\sigma^+) \frac{4\pi}{6} \sigma^3 \frac{\partial T}{\partial x}. \quad (41)$$

Let us consider the heat flux in a system at rest and of uniform density. According to Equation (6)

$$\begin{aligned} \frac{3}{2} k_B n(\mathbf{r}, t) \partial_t T(\mathbf{r}, t) &= \left\{ \frac{5}{2} \frac{1}{m\omega_0} n(\mathbf{r}, t) k_B^2 T \right. \\ &\times \left[1 + n g_2(\sigma^+) \frac{4\pi}{15} \sigma^3 \right] \\ &\left. + \frac{2}{3} \sqrt{m\pi k_B T} g_2(\sigma^+) n^2 \sigma^4 \frac{k_B}{m} \right\} \frac{\partial^2 T}{\partial x^2} \end{aligned} \quad (42)$$

yielding the thermal conductivity, λ , as the sum of the three terms inside the parenthesis in the right-hand side of Equation (42). These are the kinetic, cross-over and collisional contributions to λ , respectively. For the sake of comparison, the Enskog formulae [42] are

$$\eta_E = \eta^{(0)} [1.016/g_2(\sigma^+) + 0.8bn + 0.761g_2(\sigma^+)b^2n^2] \quad (43)$$

and

$$\lambda_E = \lambda^{(0)} [1.025/g_2(\sigma^+) + 1.230bn + 0.7764g_2(\sigma^+)b^2n^2], \quad (44)$$

where $b = 2\pi/3\sigma^3$ and $\eta^{(0)} = \frac{5}{16\sigma^2} \sqrt{\frac{mk_B T}{\pi}}$ and $\lambda^{(0)} = \frac{75}{64\sigma^2} \sqrt{\frac{k_B^3 T}{m\pi}}$.

We shall derive such corrections and compute numerically their effect by using the SMDB approach.

5. Numerical results

By using the results derived above, with the effective fields given by Equations (26), (27), (28) and (29), we construct the corresponding Lattice Boltzmann algorithm to solve the kinetic Equation (18) on a three-dimensional cubic lattice with lattice constant a , assumed to be a fraction of the hard sphere diameter σ .

The continuous dependence of $f(\mathbf{r}, \mathbf{v}, t)$ on its argument \mathbf{r} is replaced by a finite set of nodes of coordinates \mathbf{r}_i of a regular lattice spanning the physical space available to the particles. The \mathbf{v} dependence is given by spanning f over an orthonormal basis formed by a finite number of tensorial Hermite polynomials. We consider the so-called D3Q19 version, where the \mathbf{v} -space is characterised by 19 discrete velocities. The state of the system is specified by the discrete set of distribution functions $f_\alpha(\mathbf{r}_i, t)$ and for the sake of simplicity, we only consider isothermal systems and neglect the evolution of the T field. The finer details of the method have been described in Refs. [16] and [17] and will not be repeated here. In the present implementation, the method only contains the two additional terms proportional to A_{ij} and B_i .

The LB algorithm performs the following basic operations for every mesh node:

- (1) the discretised distribution is initialised by specifying the local values of the density and fluid velocity;
- (2) the values of the fields A_{ij} , B_i , C_i , Q are computed via Equations (26)–(29) using the fields \mathbf{u} and n ;
- (3) the distribution is evolved in time though the collision and streaming steps via an explicit trapezoidal rule;
- (4) the new fields $n(\mathbf{r}, t)$ and $\mathbf{u}(\mathbf{r}, t)$ are computed using the evolved distribution function;
- (5) the cycle is repeated starting from step (2).

In this scheme, one relevant question pertains the choice of the BGK relaxation frequency ω_0 . According to the Enskog picture, the kinetic contribution to viscosity should be set equal to $\eta_K = \frac{5}{16\sigma^2} \sqrt{\frac{mk_B T}{\pi}}$ and therefore $\omega_0 = \frac{nk_B T}{\eta_K}$. In addition, the low-density regime should be characterised by a constant dynamic viscosity, as first observed by Maxwell and, as $n \rightarrow 0$, the viscosity should go to zero. This rather large baseline for viscosity necessarily implies a strong departure from the local equilibrium. Such a condition is numerically critical, as it corresponds to a mean free path of the order of σ , whereas numerical stability imposes that the mean free path should be of the order of the mesh spacing a . In this case, in fact, the BGK term becomes effective in tethering the distribution close to the local equilibrium form, coherently with the small-Knudsen conditions employed in the theory developed so far. For this reason, we choose a smaller value of the kinetic viscosity. It is important to remember that also the cross-over term depends on ω_0 , but nevertheless we can still determine its effect on the global viscosity.

In the numerical benchmark, we evaluate the effect of the cross-over contributions to shear viscosity stemming from Equation (15). In this test, we choose a hard-sphere diameter $\sigma = 8a$ and the packing fraction $\chi = \frac{\pi\sigma^3 n}{6}$ ranges from dilute gas to dense liquid conditions ($\chi \simeq 0.40$). In Figure 1, we report the shear viscosity in bulk conditions as a function of the packing fraction χ . The measurements were performed by initially perturbing a uniform quiescent state by a sine wave of the type $u_z^0 \sin(q_x x)$ and monitored that its relaxation occurred with a characteristic time




Figure 1. Shear viscosity computed in bulk conditions η_{bulk} as a function of the packing fraction (this quantity should not be confused with the bulk viscosity of Section 4.1). Circles and squares are with and without the cross-over correction to viscosity, respectively.




Figure 2. Normalised shear viscosity as a function of the effective slab width ($W - \sigma$) expressed in lattice units. Circles and squares correspond to packing fractions of 0.11 and 0.32, respectively.

$\tau(q_x) = \frac{\eta}{n_0} q_x^2$, where η is the shear viscosity and n_0 the density of the uniform fluid. The data are overall smooth up to the highest packing fraction and we observe that the correction to viscosity is relevant. The presence of the crossover term determines a larger viscosity over the entire range of packing fractions studied with respect to the one obtained in the absence of it.

Next, we address the classical question regarding how the confinement in a narrow space affects the viscosity of the fluid [43–45]. In our numerical tests, we consider the flow of a hard-sphere fluid between two parallel plates and observe the density and velocity profile. Since this is a direct test of the viscosity of the fluid we expect a change with respect to the previous version where the terms A_{ij} and B_i were not taken into account. As shown in Figure 2, the measured viscosity converges for quite large channels of the order of 200 lattice units towards the predicted bulk value, whereas for small values of the width W is consistently lower, thus indicating that the walls strongly affect the motion of the adjacent layers.

6. Conclusions

In this study, we have considered the corrections to the transport coefficients which determine the cross terms between kinetic and collisional contributions. At the price of including a couple of extra contributions in the term describing the relaxation of fast modes, that is by lifting the BGK approximation for these modes, we have obtained a new estimate for the transport properties. After deriving an explicit expression for these quantities, we have evaluated the correction to viscosity via numerical simulations. It is

interesting to remark that the range of validity of the presented LB algorithm does not extend to very low densities. The reason being that, if one assumes a constant relaxation frequency ω_0 , the value of the kinematic viscosity is too small and the small Knudsen number assumption breaks down.

The resulting form of the expression for the viscosity has similarities with the one derived within the local average density model (LADM) by Davis and co-workers [46]. They assumed that the functional dependence of transport coefficients on density can be computed via the corresponding expressions in the Enskog theory of hard-sphere fluids, by replacing the actual density $n(\mathbf{r}, t)$ by the coarse grained density, $\bar{n}(\mathbf{r}, t)$. An approximation very similar to the LADM can be derived within our method by using the explicit form of Ω . Such a recipe is in the same spirit as the early weighted density density functional theories of equilibrium systems where the local free energy density was assumed to be a function of $\bar{n}(\mathbf{r})$. Moreover, as pointed out by Hoang and Galliero in a molecular dynamics study [47], the LADM underestimates the importance of the local density as far as the kinetic term is concerned. The present theory, does not suffer from such a problem since it treats on a separate basis kinetic and potential terms.

Finally, the presence of attractive interactions can be accommodated in a mean field fashion by adding a random phase term in the evolution equation. However, while representing a very important contribution to the equation of state, such a modification does not modify the transport coefficients (with the notable exception of the interspecies diffusion coefficient in mixtures, see Refs. [48] and [27]). The reason for this shortcoming is that in the Enskog-like equation the details of the potential enter only through the cross-section.

Acknowledgements

UMBM acknowledges the support of the Project COFIN-MIUR 2009.

References

- [1] T.M. Squires and S.R. Quake, *Rev. Mod. Phys.* **77**, 977 (2005).
- [2] R.B. Schoch, J. Han, and P. Renaud, *Rev. Mod. Phys.* **80**, 839 (2008).
- [3] H. Bruus, *Theoretical Microfluidics* (Oxford University Press, New York, 2008).
- [4] B.J. Kirby, *Micro and Nanoscale Fluid Mechanics* (Cambridge University Press, New York, 2010).
- [5] L. Bocquet and E. Charlaix, *Chem. Soc. Rev.* **39**, 1073 (2009).
- [6] S.K. Bhatia, M.R. Bonilla, and D. Nicholson, *Phys. Chem. Chem. Phys.* **13**, 15350 (2011).
- [7] G. Karniadakis, A. Beskok, and N. Aluru, *Microflows and Nanoflows* (Springer, New York, 2005).
- [8] L. Bocquet and J.L. Barrat, *Phys. Rev. E* **49**, 3079 (1994).
- [9] R. Hartkamp, A. Ghosh, T. Weinhart, and S. Luding, *J. Chem. Phys.* **137**, 044711 (2012).
- [10] L.A. Pohzar and K.E. Gubbins, *J. Chem. Phys.* **99**, 8970 (1993).
- [11] J.G. Anero and P. Español, *Europhys. Lett.* **78**, 50005 (2007).
- [12] A.J. Archer, *J. Phys.: Condens. Matter* **18**, 5617 (2006).
- [13] U. Marini Bettolo Marconi and S. Melchionna, *J. Chem. Phys.* **126**, 184109 (2007).
- [14] S. Succi, *The Lattice Boltzmann Equation for Fluid Dynamics and Beyond*, 1sted. (Oxford University Press, Oxford, 2001).
- [15] R. Benzi, S. Succi, and M. Vergassola, *Phys. Rep.* **222**, 145 (1992).
- [16] S. Melchionna and U. Marini Bettolo Marconi, *Europhys. Lett.* **81**, 34001 (2008).
- [17] U. Marini Bettolo Marconi and S. Melchionna, *J. Chem. Phys.* **131**, 014105 (2009).
- [18] U. Marini Bettolo Marconi, *Mol. Phys.* **109**, 1265 (2011).
- [19] L.-S. Luo, *Phys. Rev. Lett.* **62**, 1618 (1998).
- [20] P. Tarazona, *Phys. Rev. A* **31**, 2672 (1985).
- [21] R. Evans, *Fundamentals of Inhomogeneous Fluids*, edited by D. Henderson (Dekker, New York, 1992).
- [22] W. Dufty, A. Santos, and J. Brey, *Phys. Rev. Lett.* **77**, 1270 (1996).
- [23] H. van Beijeren and M.H. Ernst, *Physica A* **68**, 437 (1973); **70**, 225 (1973).
- [24] P.L. Bhatnagar, E.P. Gross, and M. Krook, *Phys. Rev.* **94**, 511 (1954).
- [25] U. Marini Bettolo Marconi and S. Melchionna, *J. Phys.: Condens. Matter* **22**, 364110 (2010).
- [26] U. Marini Bettolo Marconi and S. Melchionna, *J. Chem. Phys.* **134**, 064118 (2011).
- [27] U. Marini Bettolo Marconi and S. Melchionna, *J. Chem. Phys.* **135**, 044104 (2011).
- [28] A. Santos, J.M. Montanero, J.W. Dufty, and J.J. Brey, *Phys. Rev. E* **57**, 1644 (1998).
- [29] H.J. Kreuzer, *Non-Equilibrium Thermodynamics and Its Statistical Foundations* (Oxford University Press, New York, 1981).
- [30] S.R. de Groot and P. Mazur, *Non-Equilibrium Thermodynamics* (Dover, New York, 1984).
- [31] J.A. McLennan, *Introduction to Non-Equilibrium Statistical Mechanics* (Prentice Hall, Indiana, 1988).
- [32] G.K. Batchelor, *An Introduction to Fluid Dynamics* (Cambridge University Press, Cambridge, 1967).
- [33] L.-S. Luo, *Phys. Rev. E* **81**, 4982 (2000).
- [34] X. Shan and H. Chen, *Phys. Rev. E* **47**, 1815 (1993).
- [35] M.R. Swift, W.R. Osborn, and J.M. Yeomans, *Phys. Rev. Lett.* **75**, 830 (1995).
- [36] U. Marini Bettolo Marconi and P. Tarazona, *J. Chem. Phys.* **110**, 8032 (1999); *J. Phys.: Condens. Matter* **12**, 413 (2000).
- [37] J. Fischer and M. Methfessel, *Phys. Rev. A* **22**, 2836 (1980).
- [38] S. Melchionna and U. Marini Bettolo Marconi, *Europhys. Lett* **95**, 44002 (2011).
- [39] S. Melchionna and U. Marini Bettolo Marconi, *Phys. Rev. E* **85**, 036707 (2012).
- [40] H.C. Longuet-Higgins and J.A. Pople, *J. Chem. Phys.* **25**, 884 (1956).
- [41] J.-P. Hansen and I.R. McDonald, *Theory of Simple Liquids*, 3rded. (Academic Press, London, 2006).
- [42] S. Chapman and T.G. Cowling, *The Mathematical Theory of Non-Uniform Gases*, 3rded. (Cambridge University Press, Cambridge, 1970).

- [43] P. Travis and K.E. Gubbins, *J. Chem. Phys.* **112**, 1984 (2000).
- [44] D. Todd, J.S. Hansen, and P.J. Daivis, *Phys. Rev. Lett.* **100**, 195901 (2008).
- [45] D. Todd and J.S. Hansen, *Phys. Rev. E* **78**, 051202 (2008).
- [46] I. Bitsanis, T.K. Vanderlick, M. Tirrell, and H.T. Davis, *J. Chem. Phys.* **89**, 3152 (1988).
- [47] H. Hoang and G. Galliero, *J. Chem. Phys.* **136**, 124902 (2012).
- [48] J. Karkheck, E. Martina, and G. Stell, *Phys. Rev. A* **25**, 3328 (1982).